CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 96, 2003

A

Agostini De Manero, E., 65
Andow, D. A., 799
Archibald, S. B., 171
Arciniega, O., 144
Assael, F., 127
Azar, D., 117

B

Baselga, A., 625, 689
Bauce, E., 231
Ben-Yehuda, S., 127
Biron, D. G., 245
Bishop, D. B., 202
Boivin, G., 245
Borden, J. H., 217
Botto, E. E., 65
Boyd, D. W., Jr., 667
Bristow, C. M., 202
Britten, H. B., 856
Broeckling, C. D., 153
Broomell, B. P., 920
Brown, J. K., 65, 544
Bullock, R. C., 637

C

Calcagno, G., 563
Cam, H., 39
Carroll, S. P., 135
Castrejón, F., 672
Castrejón Gómez, V. R., 672
Chapco, W., 491
Chase, J. A., 181
Chatzimanolis, S., 225
Chong, J.-H., 539
Cisneros, R., 144, 883
Cladera, J. L., 563
Coderre, D., 245
Colauto, N. B., 448
Cole, T. J., 865
Copren, K. A., 585
Corley, J., 685
Costa, J. T., 336
Coulson, R. N., 679
Curtis, A. D., 81

D

D'Adamo, P., 685
Dang, P. M., 637
Davidson, E. W., 544
Delphia, C. M., 585
Delury, L. E., 217
Díaz, E., 144, 883
Dietrich, C. H., 700, 766
Dingle, H., 135
Dively, G. P., 631
Dmitriev, D. A., 766
Dowell, F. E., 865
Downie, D. A., 723
Dunning, K., 553

E

Ekrem, T., 265
Erwin, T. L., 596
Escher, R. L., 512
Espelie, K. E., 61
Evenden, M. L., 217

F

Felgenhauer, B. E., 870
Field Working Group, 563
Flowers, R. W., 336
Follett, P. A., 631
Forister, M. L., 73
Forrest, T. G., 433
Fritz, G. N., 86
Fuester, R. W., 553
Fujisaki, K., 793

G

Galante, E., 95, 597
Garcia De Mendoza, L., 631
Getty, G. M., 828
Greenfield, M. D., 168
Gullan, P. J., 723

H

Hammond, P. C., 847
Hancock, G., 458
Harris, M. O., 250
Hauser, M., 738
Haverty, M. I., 585, 828
Haymer, D., 594
Hendrix, D. L., 544
Hlivko, J. T., 914
Hodges, A. C., 61
Hodges, G. S., 61
Holland, J. N., 519
Hurley, R. L., 403

I

Ichiki, R., 161
Ihle, D. T., 532
Irwin, M. E., 738

J

Jameson, M. L., 415
Jancovich, J. K., 544
Johnston, J. S., 679
Jones, S. C., 181
Judd, G.J.R., 217

K

Kakimoto, T., 793
Kam, M.W.Y., 12
Katayama, N., 579
Kattan, A. D., 856
Kawano, K., 503
Keller, L., 810
Krecek, J., 181
Krieger, M. J. B., 810

L

Lane, C. P., 799
Larkin, E., 211
Levine, E., 901
Lewis, V. R., 828
Litzenberger, G., 491
Loeb, M., 344
Logarzo, G. A., 834
López, S. N., 65
Lounibos, L. P., 512
Lourenço-de-Oliveira, R., 512
Lozada, M., 685
Luzón-Ortega, J. M., 560

M

MacClure, A. R., 856
Magowski, W. L., 345
Maier, C. T., 660
Makarkin, V. N., 171
Makino, S., 652
Mangold, J. R., 181
Manrique, V., 834
Marcos-García, M. A., 458
Markwick, N. P., 250
Marler, M., 135
McCreadie, J. W., 532
McInnis, D. O., 563
McPherson, J. E., 776
Meagher, R. L., 329
Meer, R. K. V., 86
Mendel, Z., 127
Menzel, F., 448
Mescher, M. C., 810
Messing, R. H., 571
Metz, M. A., 369
Michener, C. D., 595
Micó, E., 95, 597
Miller, J. C., 847
Miyatake, T., 793
Moreau, G., 231

CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 96, 2003

A

Agostini De Manero, E., 65
Andow, D. A., 799
Archibald, S. B., 171
Arciniega, O., 144
Assael, F., 127
Azar, D., 117

B

Baselga, A., 625, 689
Bauce, E., 231
Ben-Yehuda, S., 127
Biron, D. G., 245
Bishop, D. B., 202
Boivin, G., 245
Borden, J. H., 217
Botto, E. E., 65
Boyd, D. W., Jr., 667
Bristow, C. M., 202
Britten, H. B., 856
Broeckling, C. D., 153
Broomell, B. P., 920
Brown, J. K., 65, 544
Bullock, R. C., 637

C

Calcagno, G., 563
Cam, H., 39
Carroll, S. P., 135
Castrejón, F., 672
Castrejón Gómez, V. R., 672
Chapco, W., 491
Chase, J. A., 181
Chatzimanolis, S., 225
Chong, J.-H., 539
Cisneros, R., 144, 883
Cladera, J. L., 563
Coderre, D., 245
Colauto, N. B., 448
Cole, T. J., 865
Copren, K. A., 585
Corley, J., 685
Costa, J. T., 336
Coulson, R. N., 679
Curtis, A. D., 81

D

D'Adamo, P., 685
Dang, P. M., 637
Davidson, E. W., 544
Delphia, C. M., 585
Delury, L. E., 217
Díaz, E., 144, 883
Dietrich, C. H., 700, 766
Dingle, H., 135
Dively, G. P., 631
Dmitriev, D. A., 766
Dowell, F. E., 865
Downie, D. A., 723
Dunning, K., 553

E

Ekrem, T., 265
Erwin, T. L., 596
Escher, R. L., 512
Espelie, K. E., 61
Evenden, M. L., 217

F

Felgenhauer, B. E., 870
Field Working Group, 563
Flowers, R. W., 336
Follett, P. A., 631
Forister, M. L., 73
Forrest, T. G., 433
Fritz, G. N., 86
Fuester, R. W., 553
Fujisaki, K., 793

G

Galante, E., 95, 597
Garcia De Mendoza, L., 631
Getty, G. M., 828
Greenfield, M. D., 168
Gullan, P. J., 723

H

Hammond, P. C., 847
Hancock, G., 458
Harris, M. O., 250
Hauser, M., 738
Haverty, M. I., 585, 828
Haymer, D., 594
Hendrix, D. L., 544
Hlivko, J. T., 914
Hodges, A. C., 61
Hodges, G. S., 61
Holland, J. N., 519
Hurley, R. L., 403

I

Ichiki, R., 161
Ihle, D. T., 532
Irwin, M. E., 738

J

Jameson, M. L., 415
Jancovich, J. K., 544
Johnston, J. S., 679
Jones, S. C., 181
Judd, G.J.R., 217

K

Kakimoto, T., 793
Kam, M.W.Y., 12
Katayama, N., 579
Kattan, A. D., 856
Kawano, K., 503
Keller, L., 810
Krecek, J., 181
Krieger, M. J. B., 810

L

Lane, C. P., 799
Larkin, E., 211
Levine, E., 901
Lewis, V. R., 828
Litzenberger, G., 491
Loeb, M., 344
Logarzo, G. A., 834
López, S. N., 65
Lounibos, L. P., 512
Lourenço-de-Oliveira, R., 512
Lozada, M., 685
Luzón-Ortega, J. M., 560

M

MacClure, A. R., 856
Magowski, W. L., 345
Maier, C. T., 660
Makarkin, V. N., 171
Makino, S., 652
Mangold, J. R., 181
Manrique, V., 834
Marcos-García, M. A., 458
Markwick, N. P., 250
Marler, M., 135
McCreadie, J. W., 532
McInnis, D. O., 563
McPherson, J. E., 776
Meagher, R. L., 329
Meer, R. K. V., 86
Mendel, Z., 127
Menzel, F., 448
Mescher, M. C., 810
Messing, R. H., 571
Metz, M. A., 369
Michener, C. D., 595
Micó, E., 95, 597
Miller, J. C., 847
Miyatake, T., 793
Moreau, G., 231

Morón, M. A., 597
Moser, J. C., 345

N

Nagoshi, R. N., 329
Nansen, C., 524
Nel, A., 117
Nelson, L. J., 828
Nénon, J. P., 245
Néraudeau, D., 117
Nestel, D., 237
Nieto, G., 672
Novoa, F., 625, 689

O

Oetting, R. D., 539
O'Grady, P. M., 12
Okabe, K., 652
Olmstead, K. L., 856
Olson, C. A., 595
Omwega, C. O., 865
Overholt, W. A., 865

P

Panchal, T. D., 637
Patton, J. C., 679
Paucar-Cabrera, A., 415
Pellmyr, O., 716
Pérez-Bañón, C., 458
Perreira, W. D., 12
Perrichot, V., 117
Phillips, T. W., 524
Pinto, M. A., 679
Polavarapu, S., 818
Price, L. D., 834

Q

Quesada-Allué, L. A., 237

R

Rabossi, A., 237
Ram, M. S., 865
Ramaseshiah, G., 553
Ramaswamy, S. B., 865
Roderick, G. K., 631
Roehrdanz, R. L., 901
Rojas, J. C., 672
Rosell, R. C., 544
Ross, D.N.R., 847
Ross, K. G., 810
Rotheray, G., 458
Rubink, W. L., 679
Runyon, J. B., 403
Rypstra, A. L., 914

S

Salako, J. O. B., 920
Salom, S. M., 153
Sánchez, L., 144
Sánchez-Ortega, A., 560
Sappington, T. W., 1
Saunders, J. B., 54
Saux, C., 693
Schaefer, C. W., 93, 615
Schaffner, J. C., 615
Scheffrahn, R. H., 181
Segraves, K. A., 716
Shapiro, A. M., 73
Sharkey, M. J., 169
Shaw, S. R., 834
Sheppard, W. S., 679
Shibayama, M., 883
Shields, V.D.C., 920
Shima, H., 161
Shoemaker, D. D., 810
Silva-Olivares, A., 883
Simon, C., 693
Smith, J. E., 448
Solensky, M. J., 211
Solís, A., 415
Spicer, G. S., 693
Spooner, J. D., 433
Spurgeon, D. W., 1
Steffan, S. A., 723
Su, N.-Y., 181
Sublette, J. E., 265
Sublette, M. F., 265
Suh, C. P.-C., 1
Sutherland, P. W., 250
Suzuki, N., 579
Swan, K. S., 553
Swart, C. C., 870
Szalanski, A. L., 901

T

Tanaka, S., 498
Tauber, C. A., 472
Taylor, P. B., 553
Teixeira, L.A.F., 818
Teulon, D.A.J., 107
Thaenkham, U., 44
Thanaphum, S., 44
Tierno de Figueroa, J. M., 560
Tolmasky, D., 237
Torres-Jerez, I., 65
Tripet, F., 169
Tsutsumi, V., 883
Tyler, A. M., 856

V

Val, F. C., 12
Valdes, J., 672
Van Iersel, M. W., 539
Vera, M. T., 563
Vilardi, J. C., 563
Viscarret, M. M., 65
von Dohlen, C. D., 107
Voss, S. C., 776

W

Walker, T. J., 433
Waller, D. A., 81
Wang, X.-G., 571
Weathersbee, A. A., III, 637
Webb, D. W., 369
Williams, L., III, 834
Winchell, R., 135
Wuellner, C. T., 54

Y

Yong, T.-H., 643

Z

Zhu, D.-H., 498
Zumbado, M. A., 458
Zúñiga, G., 144, 883

CUMULATIVE SUBJECT INDEX
Annals of the Entomological Society of America
Volume 96, 2003

A

A. m. scutellata,
Africanized honey bee, 679
mitochondrial DNA, 679
mitotype, 679
mtDNA haplotype, 679
Acari,
fungi, 345
host ranges, 345
phoresy, 345
Scolytidae, 345
systematics, 345
acetylene-reduction,
Reticulitermes, 81
sugar, 81
Aethiessa floralis,
Cetoniinae, 95
larval biology, 95
Oxythyrea funesta, 95
Scarabaeidae, 95
Tropinota squalida, 95
Africanized honey bee,
A. m. scutellata, 679
mitochondrial DNA, 679
mitotype, 679
mtDNA haplotype, 679
aggression,
competition, 585
fighting, 585
kin recognition, 585
subterranean termites, 585
Albian,
amber, 117
France, 117
Lebanon, 117
Lower Cretaceous, 117
Psychodidae, 117
alimentary canal, 144, 544, 883
anatomy, 144
bark beetle, 144
bark beetles, 883
Bemisia argentifolii, 544
Bemisia tabaci, 544
Dendroctonus, 144, 883
histology, 144
plant virus-vector, 544
Scolytidae, 144, 883
ultrastructure, 883
allopatry,
body size, 503
character displacement, 503
genitalia size, 503
stag beetles, 503
sympatry, 503
Alydidae,
Alydinae, 615
Apidaurus, 615
genital capsule, 615
Hemiptera, 615
Heteroptera, 615
Alydinae,
Alydidae, 615
Apidaurus, 615
genital capsule, 615
Hemiptera, 615
Heteroptera, 615
amber,
Albian, 117
France, 117
Lebanon, 117
Lower Cretaceous, 117
Psychodidae, 117
ambush predator,
flowers, 643
nectar-feeding, 643
omnivory, 643
zoophytophagy, 643
Ammonaios,
new species, 738
pupal morphology, 738
Therevidae, 738
anatomy,
alimentary canal, 144
bark beetle, 144
Dendroctonus, 144
histology, 144
Scolytidae, 144
Anisoptera,
maximum likelihood, 693
mtDNA, 693
Odonata, 693
Zygoptera, 693
Anomala, 415, 597
Anomalini, 597
Callistethus, 597
Exomala, 415
larvae, 597
larval biology, 597
morphology, 597
Phyllopertha, 415
Popillia, 415
scarab beetle, 415
Anomalini,
Anomala, 597
Callistethus, 597
larvae, 597
larval biology, 597
morphology, 597
antennal morphology,
Coccinellidae, 153
Derodontidae, 153
Laricobius nigrinus, 153
olfaction, 153
Pseudoscymnus tsugae, 153
antennal sensilla,
Lepidoptera, 672
Manilkara zapota, 672
Pyralidae, 672
Zamagiria dixolophella, 672
Anthonomus grandis,
boll weevil, 1
diapause, 1
reproductive morphology, 1
ants,
enemy-free space, 202
Homoptera, 202
mutualism, 202
aphids,
biogeography, 107
Gondwana, 107
molecular phylogeny, 107
New Zealand, 107
Tertiary origins, 107
Aphis craccivora,
extrafloral nectaries, 579
honeydew, 579
Lasius niger, 579
Tetramorium caespitum, 579
Apidaurus,
Alydidae, 615
Alydinae, 615
genital capsule, 615
Hemiptera, 615
Heteroptera, 615
apple,
Archips fuscocupreanus, 660
exotic pest, 660
host plants, 660
seasonal flight activity, 660
Tortricidae, 660
Archips fuscocupreanus,
apple, 660
exotic pest, 660
host plants, 660
seasonal flight activity, 660
Tortricidae, 660
aversion learning,
Bacillus thuringiensis, 231
behavioral response, 231
Choristoneura fumiferana, 231
dose ingestion, 231
feeding abstention, 231

B

Bacillus thuringiensis,
aversion learning, 231
behavioral response, 231
Choristoneura fumiferana, 231
dose ingestion, 231
feeding abstention, 231
Bactrocera tau,
fruit flies, 44
heat shock protein 70 cognate gene, 44
molecular marker, 44
molecular systematics, 44
species complex, 44
bark beetles,
alimentary canal, 144, 883
anatomy, 144
Dendroctonus, 144, 883
histology, 144
Scolytidae, 144, 883
ultrastructure, 883
behavior,
copulatory courtship, 225
Nordus fungicola, 225
Staphylinidae, 225
behavioral response,
aversion learning, 231
Bacillus thuringiensis, 231
Choristoneura fumiferana, 231
dose ingestion, 231
feeding abstention, 231
Belostoma lutarium,
extra-oral digestion, 870
Hemiptera, 870
mouthparts, 870
salivary gland, 870
Bemisia argentifolii,
alimentary canal, 544
Bemisia tabaci, 544
plant virus-vector, 544
Bemisia tabaci,
alimentary canal, 544
Bemisia argentifolii, 544
plant virus-vector, 544
Bemisia tabaci complex,
haplotype, 65
mitochondria COI gene, 65
phylogenetic analysis, 65
whitefly biology, 65
biocontrol,
circadian, 54
circannual phenology, 54
interspecific competition, 54
phorid parasitoids, 54
Pseudacteon, 54
biodiversity,
forest, 847
Plusiinae, 847
species richness, 847
subalpine meadows, 847
biogeography, 107, 181
aphids, 107
Caribbean, 181
Gondwana, 107
identification key, 181
molecular phylogeny, 107
new species, 181
New Zealand, 107
survey, 181
Tertiary origins, 107
biological control,
number of copulations, 553
postmating rest period, 553
reproduction, 553
sex ratio, 553
temperature at mating, 553
biotype,
chorionic egg ultrastructure, 245
Delia antiqua, 245
Delia floralis, 245
Delia radicum, 245
blueberry maggot,
founder effect, 818
highbush blueberry, 818
phenology, 818
seasonal asynchrony, 818
body size,
allopatry, 503
character displacement, 503
genitalia size, 503
stag beetles, 503
sympatry, 503
boll weevil,
Anthonomus grandis, 1
diapause, 1
reproductive morphology, 1
Bradysia difformis Frey,
Bradysia ocellaris (Comstock), 448
descriptions, 448
Diptera, 448
new synonym, 448
sciaridae, 448
Bradysia ocellaris (Comstock),
Bradysia difformis Frey, 448
descriptions, 448
Diptera, 448
new synonym, 448
sciaridae, 448
Brazil,
day length, 512
egg hatch, 512
mosquito, 512
USA, 512
Buprestidae *Prunus*,
Capnodis carbonaria, 127
Capnodis tenebrionis, 127
host selection, 127
resistance, 127
rootstock, 127

C

calling songs,
female phonoresponse, 433
katydids, 433
Phaneropterinae, 433
sibling species, 433
species concepts, 433
Callistethus,
Anomala, 597
Anomalini, 597
larvae, 597
larval biology, 597
morphology, 597
Capnodis carbonaria,
Buprestidae *Prunus*, 127
Capnodis tenebrionis, 127
host selection, 127
resistance, 127
rootstock, 127
Capnodis tenebrionis,
Buprestidae *Prunus*, 127
Capnodis carbonaria, 127
host selection, 127
resistance, 127
rootstock, 127
carbohydrates,
glutamate dehydrogenase, 237
lipids, 237
medfly, 237
metamorphosis, 237
proteins, 237
Caribbean,
biogeography, 181
identification key, 181
new species, 181
survey, 181
catalog,
Dolichopodidae, 403
key, 403
Nepalomyia, 403
Neurigonella, 403
taxonomic revision, 403
Ceraeochrysa,
Chrysopidae, 472
Chrysopodes, 472
green lacewings, 472
larvae, 472
Cetoniinae,
Aethiessa floralis, 95
larval biology, 95
Oxythyrea funesta, 95
Scarabaeidae, 95
Tropinota squalida, 95
Chalcoela pegasalis Walker,
Elasmus polistis Burks, 61
Pachysomoides fulvus Cresson, 61
Polistes metricus Say, 61
Xenos peckii Kirby, 61

character displacement,
allopatry, 503
body size, 503
genitalia size, 503
stag beetles, 503
sympatry, 503
Chelymorpha cassidea,
genetic structure, 856
habitat fragmentation, 856
heterozygosity, 856
Physonota helianthi, 856
chemical ecology,
host selection, 524
Indianmeal moth, 524
primary attraction, 524
stored-products, 524
chemotaxonomy,
colony founding, 828
Reticulitermes hesperus, 828
speciation, 828
subterranean termite, 828
chorionic egg ultrastructure,
biotype, 245
Delia antiqua, 245
Delia floralis, 245
Delia radicum, 245
Choristoneura fumiferana,
aversion learning, 231
Bacillus thuringiensis, 231
behavioral response, 231
dose ingestion, 231
feeding abstention, 231
Chroristoneura rosaceana,
ductus ejaculatorius, 217
mating status, 217
spermatophore, 217
Tortricidae, 217
Chrysomelidae,
Iberian peninsula, 689
Psylliodes, 689
taxonomy, 689
Chrysopidae,
Ceraeochrysa, 472
Chrysopodes, 472
green lacewings, 472
larvae, 472
Chrysopodes,
Ceraeochrysa, 472
Chrysopidae, 472
green lacewings, 472
larvae, 472
circadian,
biocontrol, 54
circannual phenology, 54
interspecific competition, 54
phorid parasitoids, 54
Pseudacteon, 54
circannual phenology,
biocontrol, 54
circadian, 54
interspecific competition, 54
phorid parasitoids, 54
Pseudacteon, 54
classification,
identification, 700
leafhopper, 700
morphology, 700
nomenclature, 700
cleptoparasitism,
Ensliniellinae, 652
parasitism, 652
phoresy, 652
scavenger, 652
Coccinellidae,
antennal morphology, 153
Derodontidae, 153
Laricobius nigrinus, 153
olfaction, 153
Pseudoscymnus tsugae, 153
coevolution,
Mojave desert, 716
phylogeography, 716
Prodoxidae, 716
Tegeticula, 716
colony founding,
chemotaxonomy, 828
Reticulitermes hesperus, 828
speciation, 828
subterranean termite, 828
comparative morphology,
host plants, 766
leafhopper, 766
Pinus, 766
taxonomy, 766
competition,
aggression, 585
fighting, 585
kin recognition, 585
subterranean termites, 585
Compsilura concinnata,
parasitizing site, 161
parasitoid, 161
respiration, 161
silkworm, 161
Tachinidae, 161
conservation biology,
habitat heterogeneity, 799
Lepidoptera, 799
Lycaeides melissa samuelis, 799
population dynamics, 799
copulatory courtship,
behavior, 225
Nordus fungicola, 225
Staphylinidae, 225
Cotesia flavipes,
Cotesia sesamiae, 865
near-infrared spectroscopy, 865
parasitoids, 865
species identification, 865
Cotesia sesamiae,
Cotesia flavipes, 865
near-infrared spectroscopy, 865
parasitoids, 865
species identification, 865
cotton bollworm,
food availability, 793
intraspecific predation, 793
larval development, 793
spatial pattern, 793

D

Danaus plexippus,
larval coloration, 211
melanism, 211
phenotypic plasticity, 211
polyphenism, 211
thermoregulation, 211
day length,
Brazil, 512
egg hatch, 512
mosquito, 512
USA, 512
Delia antiqua,
biotype, 245
chorionic egg ultrastructure, 245
Delia floralis, 245
Delia radicum, 245
Delia floralis,
biotype, 245
chorionic egg ultrastructure, 245
Delia antiqua, 245
Delia radicum, 245
Delia radicum,
biotype, 245
chorionic egg ultrastructure, 245
Delia antiqua, 245
Delia floralis, 245
Dendroctonus, 144, 883
alimentary canal, 144, 883
anatomy, 144
bark beetle, 144
bark beetles, 883
histology, 144
Scolytidae, 144, 883
ultrastructure, 883
Deraeocoris nigritulus,
digestive enzymes, 667
Miridae, 667
predator, 667
stylet morphology, 667
Derodontidae,
antennal morphology, 153
Coccinellidae, 153
Laricobius nigrinus, 153
olfaction, 153
Pseudoscymnus tsugae, 153
descriptions, 448, 776
Bradysia difformis Frey, 448
Bradysia ocellaris (Comstock), 448
Diptera, 448
laboratory rearing, 776

life history, 776
new synonym, 448
nymphs, 776
sciaridae, 448
Sinea diadema, 776
development,
Phenacoccus madeirensis, 539
Pseudococcidae, 539
reproduction, 539
survival, 539
Dialineura,
Incoxoverpa, 369
Nearctic revision, 369
Pallicephala, 369
diapause,
Anthonomus grandis, 1
boll weevil, 1
reproductive morphology, 1
Diaprepes abbreviatus,
18S rDNA, 637
genotyping, 637
molecular detection, 637
Pachnaeus litus, 637
digestive enzymes,
Deraeocoris nigritulus, 667
Miridae, 667
predator, 667
stylet morphology, 667
Diptera, 265, 448
Bradysia difformis Frey, 448
Bradysia ocellaris (Comstock), 448
descriptions, 448
new synonym, 448
non-biting midges, 265
North America, 265
sciaridae, 448
Tanytarsini, 265
Tanytarsus, 265
distribution,
polygyne, 86
red imported fire ant, 86
Solenopsis invicta, 86
sympatry, 86
Dolichopodidae,
catalog, 403
key, 403
Nepalomyia, 403
Neurigonella, 403
taxonomic revision, 403
dose ingestion,
aversion learning, 231
Bacillus thuringiensis, 231
behavioral response, 231
Choristoneura fumiferana, 231
feeding abstention, 231
ductus ejaculatorius,
Chroristoneura rosaceana, 217
mating status, 217
spermatophore, 217
Tortricidae, 217
Dyar's rule,
life history, 519
plant-insect interaction, 519
Upiga virescens, 519

E

ectoparasitoid,
Eurytomidae, 39
galls, 39
Hymenoptera, 39
Rileya doganlari, 39
Turkey, 39
egg hatch,
Brazil, 512
day length, 512
mosquito, 512
USA, 512
egg maturation,
Fopius arisanus, 571
host stimuli, 571
parasitoids, 571
Elasmus polistis Burks,
Chalcoela pegasalis Walker, 61
Pachysomoides fulvus Cresson, 61
Polistes metricus Say, 61
Xenos peckii Kirby, 61
electrophoresis,
gene flow, 631
sperm competition, 631
enemy-free space,
ants, 202
Homoptera, 202
mutualism, 202
Ensliniellinae,
cleptoparasitism, 652
parasitism, 652
phoresy, 652
scavenger, 652
Eocene,
fossil insects, 171
Neuroptera, 171
Okanagan Highlands, 171
Polystoechotidae, 171
Epiphyas postvittana,
histology, 250
Lepidoptera, 250
midgut cells, 250
Tortricidae, 250
toxin, 250
Eurytomidae,
ectoparasitoid, 39
galls, 39
Hymenoptera, 39
Rileya doganlari, 39
Turkey, 39
Exomala,
Anomala, 415
Phyllopertha, 415
Popillia, 415
scarab beetle, 415
exotic pest,
apple, 660
Archips fuscocupreanus, 660
host plants, 660
seasonal flight activity, 660
Tortricidae, 660
extrafloral nectaries,
Aphis craccivora, 579
honeydew, 579
Lasius niger, 579
Tetramorium caespitum, 579
extra-oral digestion,
Belostoma lutarium, 870
Hemiptera, 870
mouthparts, 870
salivary gland, 870

F

feeding abstention,
aversion learning, 231
Bacillus thuringiensis, 231
behavioral response, 231
Choristoneura fumiferana, 231
dose ingestion, 231
feeding behavior,
gustation, 920
host plant, 920
Lepidoptera, 920
Lymantriidae, 920
female phonoresponse,
calling songs, 433
katydids, 433
Phaneropterinae, 433
sibling species, 433
species concepts, 433
Ferrisia,
mealybugs, 723
phylogeny, 723
Pseudococcidae, 723
taxonomy, 723
field cages,
genetic sexing strains, 563
medfly, 563
refractory period, 563
remating, 563
sterile insect technique, 563
fighting,
aggression, 585
competition, 585
kin recognition, 585
subterranean termites, 585
fire ants,
Gp-9, 810
polygyny, 810
social organization, 810
Solenopsis invicta, 810
flight,
life history, 135
population, 135
rapid evolution, 135
seed bug, 135

wing polymorphism, 135
flight period,
Plecoptera, 560
protandry, 560
sexual dimorphism, 560
Spain, 560
flowers,
ambush predator, 643
nectar-feeding, 643
omnivory, 643
zoophytophagy, 643
food availability,
cotton bollworm, 793
intraspecific predation, 793
larval development, 793
spatial pattern, 793
food web,
herbivory, 914
Japanese beetles, 914
Mexican bean beetles, 914
spiders, 914
Fopius arisanus,
egg maturation, 571
host stimuli, 571
parasitoids, 571
forest,
biodiversity, 847
Plusiinae, 847
species richness, 847
subalpine meadows, 847
fossil insects,
Eocene, 171
Neuroptera, 171
Okanagan Highlands, 171
Polystoechotidae, 171
founder effect,
blueberry maggot, 818
highbush blueberry, 818
phenology, 818
seasonal asynchrony, 818
FR sequence,
Spodoptera frugiperda, 329
strain identification, 329
France,
Albian, 117
amber, 117
Lebanon, 117
Lower Cretaceous, 117
Psychodidae, 117
fruit flies,
Bactrocera tau, 44
heat shock protein 70 cognate gene, 44
molecular marker, 44
molecular systematics, 44
species complex, 44
fungi,
Acari, 345
host ranges, 345
phoresy, 345
Scolytidae, 345
systematics, 345

G

galls,
ectoparasitoid, 39
Eurytomidae, 39
Hymenoptera, 39
Rileya doganlari, 39
Turkey, 39
gene flow,
electrophoresis, 631
sperm competition, 631
genetic sexing strains,
field cages, 563
medfly, 563
refractory period, 563
remating, 563
sterile insect technique, 563
genetic structure,
Chelymorpha cassidea, 856
habitat fragmentation, 856
heterozygosity, 856
Physonota helianthi, 856
genital capsule,
Alydidae, 615
Alydinae, 615
Apidaurus, 615
Hemiptera, 615
Heteroptera, 615
genitalia size,
allopatry, 503
body size, 503
character displacement, 503
stag beetles, 503
sympatry, 503
genotyping,
18S rDNA, 637
Diaprepes abbreviatus, 637
molecular detection, 637
Pachnaeus litus, 637
geographic variation,
morphometrics, 73
Papilio gothica, 73
Papilio zelicaon, 73
Papilionidae, 73
seasonal variation, 73
German wasp,
local enhancement, 685
olfactory cues, 685
social insect communication, 685
yellowjackets, 685
glutamate dehydrogenase,
carbohydrates, 237
lipids, 237
medfly, 237
metamorphosis, 237
proteins, 237
Gondwana,
aphids, 107
biogeography, 107
molecular phylogeny, 107
New Zealand, 107
Tertiary origins, 107
Gp-9,
fire ants, 810
polygyny, 810
social organization, 810
Solenopsis invicta, 810
grasshoppers,
melanoplines, 491
mitochondrial DNA, 491
phylogeography, 491
podismines, 491
green lacewings,
Ceraeochrysa, 472
Chrysopidae, 472
Chrysopodes, 472
larvae, 472
group foraging,
insect communication, 336
insect sociality, 336
sawflies, 336
social insects, 336
gustation,
feeding behavior, 920
host plant, 920
Lepidoptera, 920
Lymantriidae, 920

H

habitat fragmentation,
Chelymorpha cassidea, 856
genetic structure, 856
heterozygosity, 856
Physonota helianthi, 856
habitat heterogeneity,
conservation biology, 799
Lepidoptera, 799
Lycaeides melissa samuelis, 799
population dynamics, 799
Habromyia,
larva, 458
Meromacrus, 458
Palpada, 458
puparium, 458
Syrphidae, 458
haplotype,
Bemisia tabaci complex, 65
mitochondria COI gene, 65
phylogenetic analysis, 65
whitefly biology, 65
Hawaiian *Drosophila*,
modified mouthpart species group, 12
systematics, 12
heat shock protein 70 cognate gene,
Bactrocera tau, 44
fruit flies, 44
molecular marker, 44
molecular systematics, 44
species complex, 44

Hemiptera, 615, 870
Alydidae, 615
Alydinae, 615
Apidaurus, 615
Belostoma lutarium, 870
extra-oral digestion, 870
genital capsule, 615
Heteroptera, 615
mouthparts, 870
salivary gland, 870
herbivory,
food web, 914
Japanese beetles, 914
Mexican bean beetles, 914
spiders, 914
Hesperotyphlus,
Iberian peninsula, 625
Staphylinidae, 625
taxonomy, 625
Heteroptera,
Alydidae, 615
Alydinae, 615
Apidaurus, 615
genital capsule, 615
Hemiptera, 615
heterozygosity,
Chelymorpha cassidea, 856
genetic structure, 856
habitat fragmentation, 856
Physonota helianthi, 856
highbush blueberry,
blueberry maggot, 818
founder effect, 818
phenology, 818
seasonal asynchrony, 818
histology, 144, 250
alimentary canal, 144
anatomy, 144
bark beetle, 144
Dendroctonus, 144
Epiphyas postvittana, 250
Lepidoptera, 250
midgut cells, 250
Scolytidae, 144
Tortricidae, 250
toxin, 250
Homoptera,
ants, 202
enemy-free space, 202
mutualism, 202
honeydew,
Aphis craccivora, 579
extrafloral nectaries, 579
Lasius niger, 579
Tetramorium caespitum, 579
host plants, 660, 766
apple, 660
Archips fuscocupreanus, 660
comparative morphology, 766
exotic pest, 660
feeding behavior, 920
gustation, 920
leafhopper, 766
Lepidoptera, 920
Lymantriidae, 920
Pinus, 766
seasonal flight activity, 660
taxonomy, 766
Tortricidae, 660
host ranges,
Acari, 345
fungi, 345
phoresy, 345
Scolytidae, 345
systematics, 345
host selection, 127, 524
Buprestidae *Prunus*, 127
Capnodis carbonaria, 127
Capnodis tenebrionis, 127
chemical ecology, 524
Indianmeal moth, 524
primary attraction, 524
resistance, 127
rootstock, 127
stored-products, 524
host stimuli,
egg maturation, 571
Fopius arisanus, 571
parasitoids, 571
Hydrachna magniscutata,
mites, 532
Ranatra nigra, 532
spatial distribution, 532
waterscorpions, 532
Hymenoptera,
ectoparasitoid, 39
Eurytomidae, 39
galls, 39
Rileya doganlari, 39
Turkey, 39

I

Iberian peninsula, 625, 689
Chrysomelidae, 689
Hesperotyphlus, 625
Psylliodes, 689
Staphylinidae, 625
taxonomy, 625, 689
identification,
classification, 700
leafhopper, 700
morphology, 700
nomenclature, 700
identification key,
biogeography, 181
Caribbean, 181
new species, 181
survey, 181
Incoxoverpa,
Dialineura, 369
Nearctic revision, 369
Pallicephala, 369
Indianmeal moth,
chemical ecology, 524
host selection, 524
primary attraction, 524
stored-products, 524
insect communication,
group foraging, 336
insect sociality, 336
sawflies, 336
social insects, 336
insect sociality,
group foraging, 336
insect communication, 336
sawflies, 336
social insects, 336
interspecific competition,
biocontrol, 54
circadian, 54
circannual phenology, 54
phorid parasitoids, 54
Pseudacteon, 54
intraspecific predation,
cotton bollworm, 793
food availability, 793
larval development, 793
spatial pattern, 793
ITS1,
mitochondrial DNA, 901
phylogeography, 901
polymerase chain reaction-restriction fragment-length polymorphism, 901
population genetics, 901

J

Japanese beetles,
food web, 914
herbivory, 914
Mexican bean beetles, 914
spiders, 914

K

katydids,
calling songs, 433
female phonoresponse, 433
Phaneropterinae, 433
sibling species, 433
species concepts, 433
key,
catalog, 403
Dolichopodidae, 403
Nepalomyia, 403
Neurigonella, 403
taxonomic revision, 403
kin recognition,
aggression, 585
competition, 585
fighting, 585
subterranean termites, 585

L

laboratory rearing,
descriptions, 776
life history, 776
nymphs, 776
Sinea diadema, 776
Laricobius nigrinus,
antennal morphology, 153
Coccinellidae, 153
Derodontidae, 153
olfaction, 153
Pseudoscymnus tsugae, 153
larvae, 472, 597
Anomala, 597
Anomalini, 597
Callistethus, 597
Ceraeochrysa, 472
Chrysopidae, 472
Chrysopodes, 472
green lacewings, 472
Habromyia, 458
larval biology, 597
Meromacrus, 458
morphology, 597
Palpada, 458
puparium, 458
Syrphidae, 458
larval biology, 95, 597
Aethiessa floralis, 95
Anomala, 597
Anomalini, 597
Callistethus, 597
Cetoniinae, 95
larvae, 597
morphology, 597
Oxythyrea funesta, 95
Scarabaeidae, 95
Tropinota squalida, 95
larval coloration,
Danaus plexippus, 211
melanism, 211
phenotypic plasticity, 211
polyphenism, 211
thermoregulation, 211
larval development,
cotton bollworm, 793
food availability, 793
intraspecific predation, 793
spatial pattern, 793
Lasius niger,
Aphis craccivora, 579
extrafloral nectaries, 579
honeydew, 579
Tetramorium caespitum, 579
leafhopper, 700, 766
classification, 700
comparative morphology, 766
host plants, 766
identification, 700
morphology, 700
nomenclature, 700
Pinus, 766
taxonomy, 766
Lebanon,
Albian, 117
amber, 117
France, 117
Lower Cretaceous, 117
Psychodidae, 117
Leiophron argentinensis Shaw n. sp.,
Lygus hesperus, 834
Lygus lineolaris, 834
parasitoids, 834
Taylorilygus apicalis, 834
Lepidoptera, 250, 672, 799, 920
antennal sensilla, 672
conservation biology, 799
Epiphyas postvittana, 250
feeding behavior, 920
gustation, 920
habitat heterogeneity, 799
histology, 250
host plant, 920
Lycaeides melissa samuelis, 799
Lymantriidae, 920
Manilkara zapota, 672
midgut cells, 250
population dynamics, 799
Pyralidae, 672
Tortricidae, 250
toxin, 250
Zamagiria dixolophella, 672
life history, 135, 519, 776
descriptions, 776
Dyar's rule, 519
flight, 135
laboratory rearing, 776
nymphs, 776
plant-insect interaction, 519
population, 135
rapid evolution, 135
seed bug, 135
Sinea diadema, 776
Upiga virescens, 519
wing polymorphism, 135
lipids,
carbohydrates, 237
glutamate dehydrogenase, 237
medfly, 237
metamorphosis, 237
proteins, 237
local enhancement,
German wasp, 685
olfactory cues, 685
social insect communication, 685
yellowjackets, 685
Locusta migratoria,
phase polymorphism, 498
precopulatory mounting, 498
sperm competition, 498
Lower Cretaceous,
Albian, 117
amber, 117
France, 117
Lebanon, 117
Psychodidae, 117
Lycaeides melissa samuelis,
conservation biology, 799
habitat heterogeneity, 799
Lepidoptera, 799
population dynamics, 799
Lygus hesperus,
Leiophron argentinensis Shaw n. sp., 834
Lygus lineolaris, 834
parasitoids, 834
Taylorilygus apicalis, 834
Lygus lineolaris,
Leiophron argentinensis Shaw n. sp., 834
Lygus hesperus, 834
parasitoids, 834
Taylorilygus apicalis, 834
Lymantriidae,
feeding behavior, 920
gustation, 920
host plant, 920
Lepidoptera, 920

M

Manilkara zapota,
antennal sensilla, 672
Lepidoptera, 672
Pyralidae, 672
Zamagiria dixolophella, 672
mating status,
Chroristoneura rosaceana, 217
ductus ejaculatorius, 217
spermatophore, 217
Tortricidae, 217
maximum likelihood,
Anisoptera, 693
mtDNA, 693
Odonata, 693
Zygoptera, 693
mealybugs,
Ferrisia, 723
phylogeny, 723
Pseudococcidae, 723
taxonomy, 723
medfly, 237, 563
carbohydrates, 237
field cages, 563
genetic sexing strains, 563
glutamate dehydrogenase, 237
lipids, 237
metamorphosis, 237
proteins, 237
refractory period, 563
remating, 563
sterile insect technique, 563

melanism,
Danaus plexippus, 211
larval coloration, 211
phenotypic plasticity, 211
polyphenism, 211
thermoregulation, 211
melanoplines,
grasshoppers, 491
mitochondrial DNA, 491
phylogeography, 491
podismines, 491
Meromacrus,
Habromyia, 458
larva, 458
Palpada, 458
puparium, 458
Syrphidae, 458
metamorphosis,
carbohydrates, 237
glutamate dehydrogenase, 237
lipids, 237
medfly, 237
proteins, 237
Mexican bean beetles,
food web, 914
herbivory, 914
Japanese beetles, 914
spiders, 914
midgut cells,
Epiphyas postvittana, 250
histology, 250
Lepidoptera, 250
Tortricidae, 250
toxin, 250
Miridae,
Deraeocoris nigritulus, 667
digestive enzymes, 667
predator, 667
stylet morphology, 667
mites,
Hydrachna magniscutata, 532
Ranatra nigra, 532
spatial distribution, 532
waterscorpions, 532
mitochondria COI gene,
Bemisia tabaci complex, 65
haplotype, 65
phylogenetic analysis, 65
whitefly biology, 65
mitochondrial DNA, 491, 679, 901
A. m. scutellata, 679
Africanized honey bee, 679
grasshoppers, 491
ITS1, 901
melanoplines, 491
mitotype, 679
mtDNA haplotype, 679
phylogeography, 491, 901
podismines, 491
polymerase chain reaction-restriction fragment-length polymorphism, 901
population genetics, 901
mitotype,
A. m. scutellata, 679
Africanized honey bee, 679
mitochondrial DNA, 679
mtDNA haplotype, 679
modified mouthpart species group,
Hawaiian *Drosophila*, 12
systematics, 12
Mojave desert,
coevolution, 716
phylogeography, 716
Prodoxidae, 716
Tegeticula, 716
molecular detection,
18S rDNA, 637
Diaprepes abbreviatus, 637
genotyping, 637
Pachnaeus litus, 637
molecular marker,
Bactrocera tau, 44
fruit flies, 44
heat shock protein 70 cognate gene, 44
molecular systematics, 44
species complex, 44
molecular phylogeny,
aphids, 107
biogeography, 107
Gondwana, 107
New Zealand, 107
Tertiary origins, 107
molecular systematics,
Bactrocera tau, 44
fruit flies, 44
heat shock protein 70 cognate gene, 44
molecular marker, 44
species complex, 44
morphology, 597, 700
Anomala, 597
Anomalini, 597
Callistethus, 597
classification, 700
identification, 700
larvae, 597
larval biology, 597
leafhopper, 700
nomenclature, 700
morphometrics,
geographic variation, 73
Papilio gothica, 73
Papilio zelicaon, 73
Papilionidae, 73
seasonal variation, 73
mosquito,
Brazil, 512
day length, 512
egg hatch, 512
USA, 512
mouthparts,
Belostoma lutarium, 870
extra-oral digestion, 870
Hemiptera, 870
salivary gland, 870
mtDNA,
Anisoptera, 693
maximum likelihood, 693
Odonata, 693
Zygoptera, 693
mtDNA haplotype,
A. m. scutellata, 679
Africanized honey bee, 679
mitochondrial DNA, 679
mitotype, 679
mutualism,
ants, 202
enemy-free space, 202
Homoptera, 202

N

Nearctic revision,
Dialineura, 369
Incoxoverpa, 369
Pallicephala, 369
near-infrared spectroscopy,
Cotesia flavipes, 865
Cotesia sesamiae, 865
parasitoids, 865
species identification, 865
nectar-feeding,
ambush predator, 643
flowers, 643
omnivory, 643
zoophytophagy, 643
Nepalomyia,
catalog, 403
Dolichopodidae, 403
key, 403
Neurigonella, 403
taxonomic revision, 403
Neurigonella,
catalog, 403
Dolichopodidae, 403
key, 403
Nepalomyia, 403
taxonomic revision, 403
Neuroptera,
Eocene, 171
fossil insects, 171
Okanagan Highlands, 171
Polystoechotidae, 171
new species, 181, 738
Ammonaios, 738
biogeography, 181
Caribbean, 181
identification key, 181
pupal morphology, 738
survey, 181
Therevidae, 738

new synonym,
Bradysia difformis Frey, 448
Bradysia ocellaris (Comstock), 448
descriptions, 448
Diptera, 448
sciaridae, 448
New Zealand,
aphids, 107
biogeography, 107
Gondwana, 107
molecular phylogeny, 107
Tertiary origins, 107
nomenclature,
classification, 700
identification, 700
leafhopper, 700
morphology, 700
non-biting midges,
Diptera, 265
North America, 265
Tanytarsini, 265
Tanytarsus, 265
Nordus fungicola,
behavior, 225
copulatory courtship, 225
Staphylinidae, 225
North America,
Diptera, 265
non-biting midges, 265
Tanytarsini, 265
Tanytarsus, 265
number of copulations,
biological control, 553
postmating rest period, 553
reproduction, 553
sex ratio, 553
temperature at mating, 553
nymphs,
descriptions, 776
laboratory rearing, 776
life history, 776
Sinea diadema, 776

O

Odonata,
Anisoptera, 693
maximum likelihood, 693
mtDNA, 693
Zygoptera, 693
Okanagan Highlands,
Eocene, 171
fossil insects, 171
Neuroptera, 171
Polystoechotidae, 171
olfaction,
antennal morphology, 153
Coccinellidae, 153
Derodontidae, 153
Laricobius nigrinus, 153
Pseudoscymnus tsugae, 153
olfactory cues,
German wasp, 685
local enhancement, 685
social insect communication, 685
yellowjackets, 685
omnivory,
ambush predator, 643
flowers, 643
nectar-feeding, 643
zoophytophagy, 643
Oxythyrea funesta,
Aethiessa floralis, 95
Cetoniinae, 95
larval biology, 95
Scarabaeidae, 95
Tropinota squalida, 95

P

Pachnaeus litus,
18S rDNA, 637
Diaprepes abbreviatus, 637
genotyping, 637
molecular detection, 637
Pachysomoides fulvus Cresson,
Chalcoela pegasalis Walker, 61
Elasmus polistis Burks, 61
Polistes metricus Say, 61
Xenos peckii Kirby, 61
Pallicephala,
Dialineura, 369
Incoxoverpa, 369
Nearctic revision, 369
Palpada,
Habromyia, 458
larva, 458
Meromacrus, 458
puparium, 458
Syrphidae, 458
Papilio gothica,
geographic variation, 73
morphometrics, 73
Papilio zelicaon, 73
Papilionidae, 73
seasonal variation, 73
Papilio zelicaon,
geographic variation, 73
morphometrics, 73
Papilio gothica, 73
Papilionidae, 73
seasonal variation, 73
Papilionidae,
geographic variation, 73
morphometrics, 73
Papilio gothica, 73
Papilio zelicaon, 73
seasonal variation, 73
parasitism,
cleptoparasitism, 652
Ensliniellinae, 652
phoresy, 652
scavenger, 652
parasitizing site,
Compsilura concinnata, 161
parasitoid, 161
respiration, 161
silkworm, 161
Tachinidae, 161
parasitoids, 571, 834, 865
Compsilura concinnata, 161
Cotesia flavipes, 865
Cotesia sesamiae, 865
egg maturation, 571
Fopius arisanus, 571
host stimuli, 571
Leiophron argentinensis Shaw n. sp., 834
Lygus hesperus, 834
Lygus lineolaris, 834
near-infrared spectroscopy, 865
parasitizing site, 161
respiration, 161
silkworm, 161
species identification, 865
Tachinidae, 161
Taylorilygus apicalis, 834
Phaneropterinae,
calling songs, 433
female phonoresponse, 433
katydids, 433
sibling species, 433
species concepts, 433
phase polymorphism,
Locusta migratoria, 498
precopulatory mounting, 498
sperm competition, 498
Phenacoccus madeirensis,
development, 539
Pseudococcidae, 539
reproduction, 539
survival, 539
phenology,
blueberry maggot, 818
founder effect, 818
highbush blueberry, 818
seasonal asynchrony, 818
phenotypic plasticity,
Danaus plexippus, 211
larval coloration, 211
melanism, 211
polyphenism, 211
thermoregulation, 211
phoresy, 345, 652
Acari, 345
cleptoparasitism, 652
Ensliniellinae, 652
fungi, 345
host ranges, 345
parasitism, 652
scavenger, 652
Scolytidae, 345
systematics, 345

phorid parasitoids,
biocontrol, 54
circadian, 54
circannual phenology, 54
interspecific competition, 54
Pseudacteon, 54
Phyllopertha,
Anomala, 415
Exomala, 415
Popillia, 415
scarab beetle, 415
phylogenetic analysis,
Bemisia tabaci complex, 65
haplotype, 65
mitochondria COI gene, 65
whitefly biology, 65
phylogeny,
Ferrisia, 723
mealybugs, 723
Pseudococcidae, 723
taxonomy, 723
phylogeography, 491, 716, 901
coevolution, 716
grasshoppers, 491
ITS1, 901
melanoplines, 491
mitochondrial DNA, 491, 901
Mojave desert, 716
podismines, 491
polymerase chain reaction-restriction fragment-length polymorphism, 901
population genetics, 901
Prodoxidae, 716
Tegeticula, 716
Physonota helianthi,
Chelymorpha cassidea, 856
genetic structure, 856
habitat fragmentation, 856
heterozygosity, 856
Pinus,
comparative morphology, 766
host plants, 766
leafhopper, 766
taxonomy, 766
plant virus-vector,
alimentary canal, 544
Bemisia argentifolii, 544
Bemisia tabaci, 544
plant-insect interaction,
Dyar's rule, 519
life history, 519
Upiga virescens, 519
Plecoptera,
flight period, 560
protandry, 560
sexual dimorphism, 560
Spain, 560
Plusiinae,
biodiversity, 847
forest, 847
species richness, 847
subalpine meadows, 847
podismines,
grasshoppers, 491
melanoplines, 491
mitochondrial DNA, 491
phylogeography, 491
Polistes metricus Say,
Chalcoela pegasalis Walker, 61
Elasmus polistis Burks, 61
Pachysomoides fulvus Cresson, 61
Xenos peckii Kirby, 61
polygyne,
distribution, 86
red imported fire ant, 86
Solenopsis invicta, 86
sympatry, 86
polygyny,
fire ants, 810
Gp-9, 810
social organization, 810
Solenopsis invicta, 810
polymerase chain reaction-restriction fragment-length polymorphism,
ITS1, 901
mitochondrial DNA, 901
phylogeography, 901
population genetics, 901
polyphenism,
Danaus plexippus, 211
larval coloration, 211
melanism, 211
phenotypic plasticity, 211
thermoregulation, 211
Polystoechotidae,
Eocene, 171
fossil insects, 171
Neuroptera, 171
Okanagan Highlands, 171
Popillia,
Anomala, 415
Exomala, 415
Phyllopertha, 415
scarab beetle, 415
population,
flight, 135
life history, 135
rapid evolution, 135
seed bug, 135
wing polymorphism, 135
population dynamics,
conservation biology, 799
habitat heterogeneity, 799
Lepidoptera, 799
Lycaeides melissa samuelis, 799
population genetics,
ITS1, 901
mitochondrial DNA, 901
phylogeography, 901
polymerase chain reaction-restriction fragment-length polymorphism, 901
postmating rest period,
biological control, 553
number of copulations, 553
reproduction, 553
sex ratio, 553
temperature at mating, 553
precopulatory mounting,
Locusta migratoria, 498
phase polymorphism, 498
sperm competition, 498
predator,
Deraeocoris nigritulus, 667
digestive enzymes, 667
Miridae, 667
stylet morphology, 667
primary attraction,
chemical ecology, 524
host selection, 524
Indianmeal moth, 524
stored-products, 524
Prodoxidae,
coevolution, 716
Mojave desert, 716
phylogeography, 716
Tegeticula, 716
protandry,
flight period, 560
Plecoptera, 560
sexual dimorphism, 560
Spain, 560
proteins,
carbohydrates, 237
glutamate dehydrogenase, 237
lipids, 237
medfly, 237
metamorphosis, 237
Pseudacteon,
biocontrol, 54
circadian, 54
circannual phenology, 54
interspecific competition, 54
phorid parasitoids, 54
Pseudococcidae, 539, 723
development, 539
Ferrisia, 723
mealybugs, 723
Phenacoccus madeirensis, 539
phylogeny, 723
reproduction, 539
survival, 539
taxonomy, 723
Pseudoscymnus tsugae,
antennal morphology, 153
Coccinellidae, 153
Derodontidae, 153
Laricobius nigrinus, 153
olfaction, 153

Psychodidae,
Albian, 117
amber, 117
France, 117
Lebanon, 117
Lower Cretaceous, 117
Psylliodes,
Chrysomelidae, 689
Iberian peninsula, 689
taxonomy, 689
pupal morphology,
Ammonaios, 738
new species, 738
Therevidae, 738
puparium,
Habromyia, 458
larva, 458
Meromacrus, 458
Palpada, 458
Syrphidae, 458
Pyralidae,
antennal sensilla, 672
Lepidoptera, 672
Manilkara zapota, 672
Zamagiria dixolophella, 672

R

Ranatra nigra,
Hydrachna magniscutata, 532
mites, 532
spatial distribution, 532
waterscorpions, 532
rapid evolution,
flight, 135
life history, 135
population, 135
seed bug, 135
wing polymorphism, 135
18S rDNA,
Diaprepes abbreviatus, 637
genotyping, 637
molecular detection, 637
Pachnaeus litus, 637
red imported fire ant,
distribution, 86
polygyne, 86
Solenopsis invicta, 86
sympatry, 86
refractory period,
field cages, 563
genetic sexing strains, 563
medfly, 563
remating, 563
sterile insect technique, 563
remating,
field cages, 563
genetic sexing strains, 563
medfly, 563
refractory period, 563
sterile insect technique, 563
reproduction, 539, 553
biological control, 553
development, 539
number of copulations, 553
Phenacoccus madeirensis, 539
postmating rest period, 553
Pseudococcidae, 539
sex ratio, 553
survival, 539
temperature at mating, 553
reproductive morphology,
Anthonomus grandis, 1
boll weevil, 1
diapause, 1
resistance,
Buprestidae *Prunus*, 127
Capnodis carbonaria, 127
Capnodis tenebrionis, 127
host selection, 127
rootstock, 127
respiration,
Compsilura concinnata, 161
parasitizing site, 161
parasitoid, 161
silkworm, 161
Tachinidae, 161
Reticulitermes,
acetylene-reduction, 81
sugar, 81
Reticulitermes hesperus,
chemotaxonomy, 828
colony founding, 828
speciation, 828
subterranean termite, 828
Rileya doganlari,
ectoparasitoid, 39
Eurytomidae, 39
galls, 39
Hymenoptera, 39
Turkey, 39
rootstock,
Buprestidae *Prunus*, 127
Capnodis carbonaria, 127
Capnodis tenebrionis, 127
host selection, 127
resistance, 127

S

salivary gland,
Belostoma lutarium, 870
extra-oral digestion, 870
Hemiptera, 870
mouthparts, 870
sawflies,
group foraging, 336
insect communication, 336
insect sociality, 336
social insects, 336
scarab beetle,
Anomala, 415
Exomala, 415
Phyllopertha, 415
Popillia, 415
Scarabaeidae,
Aethiessa floralis, 95
Cetoniinae, 95
larval biology, 95
Oxythyrea funesta, 95
Tropinota squalida, 95
scavenger,
cleptoparasitism, 652
Ensliniellinae, 652
parasitism, 652
phoresy, 652
sciaridae,
Bradysia difformis Frey, 448
Bradysia ocellaris (Comstock), 448
descriptions, 448
Diptera, 448
new synonym, 448
Scolytidae, 144, 345, 883
Acari, 345
alimentary canal, 144, 883
anatomy, 144
bark beetle, 144
bark beetles, 883
Dendroctonus, 144, 883
fungi, 345
histology, 144
host ranges, 345
phoresy, 345
systematics, 345
ultrastructure, 883
seasonal asynchrony,
blueberry maggot, 818
founder effect, 818
highbush blueberry, 818
phenology, 818
seasonal flight activity,
apple, 660
Archips fuscocupreanus, 660
exotic pest, 660
host plants, 660
Tortricidae, 660
seasonal variation,
geographic variation, 73
morphometrics, 73
Papilio gothica, 73
Papilio zelicaon, 73
Papilionidae, 73
seed bug,
flight, 135
life history, 135
population, 135
rapid evolution, 135
wing polymorphism, 135
sex ratio,
biological control, 553
number of copulations, 553
postmating rest period, 553
reproduction, 553
temperature at mating, 553

sexual dimorphism,
flight period, 560
Plecoptera, 560
protandry, 560
Spain, 560
sibling species,
calling songs, 433
female phonoresponse, 433
katydids, 433
Phaneropterinae, 433
species concepts, 433
silkworm,
Compsilura concinnata, 161
parasitizing site, 161
parasitoid, 161
respiration, 161
Tachinidae, 161
Sinea diadema,
descriptions, 776
laboratory rearing, 776
life history, 776
nymphs, 776
social insect communication,
German wasp, 685
local enhancement, 685
olfactory cues, 685
yellowjackets, 685
social insects,
group foraging, 336
insect communication, 336
insect sociality, 336
sawflies, 336
social organization,
fire ants, 810
Gp-9, 810
polygyny, 810
Solenopsis invicta, 810
Solenopsis invicta, 86, 810
distribution, 86
fire ants, 810
Gp-9, 810
polygyne, 86
polygyny, 810
red imported fire ant, 86
social organization, 810
sympatry, 86
Spain,
flight period, 560
Plecoptera, 560
protandry, 560
sexual dimorphism, 560
spatial distribution,
Hydrachna magniscutata, 532
mites, 532
Ranatra nigra, 532
waterscorpions, 532
spatial pattern,
cotton bollworm, 793
food availability, 793
intraspecific predation, 793
larval development, 793
speciation,
chemotaxonomy, 828
colony founding, 828
Reticulitermes hesperus, 828
subterranean termite, 828
species complex,
Bactrocera tau, 44
fruit flies, 44
heat shock protein 70 cognate gene, 44
molecular marker, 44
molecular systematics, 44
species concepts,
calling songs, 433
female phonoresponse, 433
katydids, 433
Phaneropterinae, 433
sibling species, 433
species identification,
Cotesia flavipes, 865
Cotesia sesamiae, 865
near-infrared spectroscopy, 865
parasitoids, 865
species richness,
biodiversity, 847
forest, 847
Plusiinae, 847
subalpine meadows, 847
sperm competition, 498, 631
electrophoresis, 631
gene flow, 631
Locusta migratoria, 498
phase polymorphism, 498
precopulatory mounting, 498
spermatophore,
Chroristoneura rosaceana, 217
ductus ejaculatorius, 217
mating status, 217
Tortricidae, 217
spiders,
food web, 914
herbivory, 914
Japanese beetles, 914
Mexican bean beetles, 914
Spodoptera frugiperda,
FR sequence, 329
strain identification, 329
stag beetles,
allopatry, 503
body size, 503
character displacement, 503
genitalia size, 503
sympatry, 503
Staphylinidae, 225, 625
behavior, 225
copulatory courtship, 225
Hesperotyphlus, 625
Iberian peninsula, 625
Nordus fungicola, 225
taxonomy, 625
sterile insect technique,
field cages, 563
genetic sexing strains, 563
medfly, 563
refractory period, 563
remating, 563
stored-products,
chemical ecology, 524
host selection, 524
Indianmeal moth, 524
primary attraction, 524
strain identification,
FR sequence, 329
Spodoptera frugiperda, 329
stylet morphology,
Deraeocoris nigritulus, 667
digestive enzymes, 667
Miridae, 667
predator, 667
subalpine meadows,
biodiversity, 847
forest, 847
Plusiinae, 847
species richness, 847
subterranean termites,
aggression, 585
chemotaxonomy, 828
colony founding, 828
competition, 585
fighting, 585
kin recognition, 585
Reticulitermes hesperus, 828
speciation, 828
sugar,
acetylene-reduction, 81
Reticulitermes, 81
survey,
biogeography, 181
Caribbean, 181
identification key, 181
new species, 181
survival,
development, 539
Phenacoccus madeirensis, 539
Pseudococcidae, 539
reproduction, 539
sympatry, 86, 503
allopatry, 503
body size, 503
character displacement, 503
distribution, 86
genitalia size, 503
polygyne, 86
red imported fire ant, 86
Solenopsis invicta, 86
stag beetles, 503
Syrphidae,
Habromyia, 458
larva, 458
Meromacrus, 458
Palpada, 458
puparium, 458

systematics, 12, 345
Acari, 345
fungi, 345
Hawaiian *Drosophila*, 12
host ranges, 345
modified mouthpart species group, 12
phoresy, 345
Scolytidae, 345

T

Tachinidae,
Compsilura concinnata, 161
parasitizing site, 161
parasitoid, 161
respiration, 161
silkworm, 161
Tanytarsini,
Diptera, 265
non-biting midges, 265
North America, 265
Tanytarsus, 265
Tanytarsus,
Diptera, 265
non-biting midges, 265
North America, 265
Tanytarsini, 265
taxonomic revision,
catalog, 403
Dolichopodidae, 403
key, 403
Nepalomyia, 403
Neurigonella, 403
taxonomy, 625, 689, 723, 766
Chrysomelidae, 689
comparative morphology, 766
Ferrisia, 723
Hesperotyphlus, 625
host plants, 766
Iberian peninsula, 625, 689
leafhopper, 766
mealybugs, 723
phylogeny, 723
Pinus, 766
Pseudococcidae, 723
Psylliodes, 689
Staphylinidae, 625
Taylorilygus apicalis,
Leiophron argentinensis Shaw n. sp., 834
Lygus hesperus, 834
Lygus lineolaris, 834
parasitoids, 834
Tegeticula,
coevolution, 716
Mojave desert, 716
phylogeography, 716
Prodoxidae, 716
temperature at mating,
biological control, 553
number of copulations, 553
postmating rest period, 553
reproduction, 553
sex ratio, 553
Tertiary origins,
aphids, 107
biogeography, 107
Gondwana, 107
molecular phylogeny, 107
New Zealand, 107
Tetramorium caespitum,
Aphis craccivora, 579
extrafloral nectaries, 579
honeydew, 579
Lasius niger, 579
Therevidae,
Ammonaios, 738
new species, 738
pupal morphology, 738
thermoregulation,
Danaus plexippus, 211
larval coloration, 211
melanism, 211
phenotypic plasticity, 211
polyphenism, 211
Tortricidae, 217, 250, 660
apple, 660
Archips fuscocupreanus, 660
Chroristoneura rosaceana, 217
ductus ejaculatorius, 217
Epiphyas postvittana, 250
exotic pest, 660
histology, 250
host plants, 660
Lepidoptera, 250
mating status, 217
midgut cells, 250
seasonal flight activity, 660
spermatophore, 217
toxin, 250
toxin,
Epiphyas postvittana, 250
histology, 250
Lepidoptera, 250
midgut cells, 250
Tortricidae, 250
Tropinota squalida,
Aethiessa floralis, 95
Cetoniinae, 95
larval biology, 95
Oxythyrea funesta, 95
Scarabaeidae, 95
Turkey,
ectoparasitoid, 39
Eurytomidae, 39
galls, 39
Hymenoptera, 39
Rileya doganlari, 39

U

ultrastructure,
alimentary canal, 883
bark beetles, 883
Dendroctonus, 883
Scolytidae, 883
Upiga virescens,
Dyar's rule, 519
life history, 519
plant-insect interaction, 519
USA,
Brazil, 512
day length, 512
egg hatch, 512
mosquito, 512

W

waterscorpions,
Hydrachna magniscutata, 532
mites, 532
Ranatra nigra, 532
spatial distribution, 532
whitefly biology,
Bemisia tabaci complex, 65
haplotype, 65
mitochondria COI gene, 65
phylogenetic analysis, 65
wing polymorphism,
flight, 135
life history, 135
population, 135
rapid evolution, 135
seed bug, 135

X

Xenos peckii Kirby,
Chalcoela pegasalis Walker, 61
Elasmus polistis Burks, 61
Pachysomoides fulvus Cresson, 61
Polistes metricus Say, 61

Y

yellowjackets,
German wasp, 685
local enhancement, 685
olfactory cues, 685
social insect communication, 685

Z

Zamagiria dixolophella,
antennal sensilla, 672
Lepidoptera, 672
Manilkara zapota, 672
Pyralidae, 672
zoophytophagy,
ambush predator, 643
flowers, 643
nectar-feeding, 643
omnivory, 643
Zygoptera,
Anisoptera, 693
maximum likelihood, 693
mtDNA, 693
Odonata, 693